

United States Patent and Trademark Office

UNITED STATES DEPARTMENT OF COMMERCE United States Patent and Trademark Office Address: COMMISSIONER FOR PATENTS P.O. Box 1450 Alexandria, Virginia 22313-1450 www.uspto.gov

DATE MAILED: 06/27/2006

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
10/666,535	09/22/2003	Hideki Ichikawa	2923-0562	5882
6449 7	590 06/27/2006	EXAMINER		
	, FIGG, ERNST & MA	ROMEO, DAVID S		
1425 K STREET, N.W. SUITE 800			ART UNIT	PAPER NUMBER
	N, DC 20005		1647	

Please find below and/or attached an Office communication concerning this application or proceeding.

	Application No.	Applicant(s)			
	10/666,535	ICHIKAWA ET AL.			
Office Action Summary	Examiner	Art Unit			
	David S. Romeo	1647			
The MAILING DATE of this communication a Period for Reply	ppears on the cover sheet wit	th the correspondence address			
A SHORTENED STATUTORY PERIOD FOR REF WHICHEVER IS LONGER, FROM THE MAILING - Extensions of time may be available under the provisions of 37 CFR after SIX (6) MONTHS from the mailing date of this communication. - If NO period for reply is specified above, the maximum statutory period - Failure to reply within the set or extended period for reply will, by stat Any reply received by the Office later than three months after the mail earned patent term adjustment. See 37 CFR 1.704(b).	DATE OF THIS COMMUNIC 1.136(a). In no event, however, may a re od will apply and will expire SIX (6) MON ute, cause the application to become AB	CATION. THS from the mailing date of this communication. ANDONED (35 U.S.C. § 133).			
Status					
1) Responsive to communication(s) filed on 22	September 2003.				
	nis action is non-final.				
3)☐ Since this application is in condition for allow					
closed in accordance with the practice under	r <i>Ex parte Quayle</i> , 1935 C.D.	. 11, 453 O.G. 213.			
Disposition of Claims					
4)⊠ Claim(s) <u>1-14</u> is/are pending in the application	on.				
4a) Of the above claim(s) is/are withdo		•			
5) Claim(s) is/are allowed.					
6)⊠ Claim(s) <u>1-14</u> is/are rejected.					
7) Claim(s) is/are objected to.					
8) Claim(s) are subject to restriction and	or election requirement.				
Application Papers		•			
9) The specification is objected to by the Exami	ner.				
10)☐ The drawing(s) filed on is/are: a)☐ ac		ov the Examiner.			
Applicant may not request that any objection to the					
Replacement drawing sheet(s) including the соле		· ·			
11) The oath or declaration is objected to by the					
Priority under 35 U.S.C. § 119					
12)⊠ Acknowledgment is made of a claim for foreig	gn priority under 35 U.S.C. §	119(a)-(d) or (f).			
a)⊠ All b)□ Some * c)□ None of:					
1. Certified copies of the priority docume	nts have been received.				
2. Certified copies of the priority docume	nts have been received in Ap	oplication No. <u>09/355,551</u> .			
3. Copies of the certified copies of the pr	iority documents have been	received in this National Stage			
application from the International Bure	au (PCT Rule 17.2(a)).				
* See the attached detailed Office action for a list	st of the certified copies not r	received.			
Attachment(s)					
1) Notice of References Cited (PTO-892)	4) Interview St	ummary (PTO-413)			
2) Notice of Draftsperson's Patent Drawing Review (PTO-948)	Paper No(s)	/Mail Date			
3) Information Disclosure Statement(s) (PTO-1449 or PTO/SB/0-Paper No(s)/Mail Date <u>0903</u> .	8)	formal Patent Application (PTO-152) —·			
J.S. Patent and Trademark Office					

Art Unit: 1647

15

20

25

DETAILED ACTION

Page 2

Claims 1–14 are pending and being examined.

Priority

Under 35 U.S.C. 120, the claims in a U.S. application are entitled to the benefit of the filing date of an earlier filed U.S. application if the subject matter of the claim is disclosed in the manner provided by 35 U.S.C. 112, first paragraph in the earlier filed application. Under 35 U.S.C. 119 (a) or (e), the claims in a U.S. application are entitled to the benefit of a foreign priority date or the filing date of a provisional application if the corresponding foreign application or provisional application supports the claims in the manner required by 35 U.S.C. 112, first paragraph.

In the present case the subject matter of claims 8–11 was not disclosed in the manner provided by 35 U.S.C. 112, first paragraph in the earlier filed 09/355,551 application, and the corresponding foreign application does not support claims 8–11 in the manner required by 35 U.S.C. 112, first paragraph. Accordingly, the effective filing date of claims 8–11 is 09/22/2003.

Claim Rejections - 35 USC § 103

The following is a quotation of 35 U.S.C. 103(a) which forms the basis for all obviousness rejections set forth in this Office action:

(a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negatived by the manner in which the invention was made.

Claims 1–7 and 12–14 are rejected under 35 U.S.C. 103(a) as being unpatentable over Neidhardt (WO 93/16099) in view of Ron (U. S. Patent No. 5,171,579) and Avis (1990).

Neidhardt teaches osteogenic protein MP52 (see SEQ ID NOs: 1 and 3), the recombinant

Art Unit: 1647

5

10

15

20

production thereof (paragraph bridging pages 6-7 through paragraph bridging pages 7-8), and a pharmaceutical composition comprising same for bone healing (page 9, full paragraph 1). The limitation "produced by a means of genetic engineering technology" is a product-by-process limitation. If the MP52 produced by the process is the same as or obvious from Neidhardt's MP52, the MP52 is unpatentable even though it may have been made by a different process because the process limitation is not viewed as positively limiting the MP52, as it is assumed that equivalent products are obtainable by multiple routes. The burden is upon the applicants to establish a patentable distinction between the MP52 produced by the process and Neidhardt's MP52. Neidhardt does not teach lyophilization of MP52 in the presence of mannitol.

Ron teaches that osteogenic proteins can be utilized in the form of a pharmaceutically acceptable solution (including reconstitution from a lyophilized form). It is optimal to solubilize the osteogenic protein at concentrations of at least about 2 mg/ml, preferably about 4 mg/ml, so that a pharmaceutically effective amount of protein can be delivered without undue volumes of carrier being necessary (column 2, lines 22-29). Ron teaches that additional optional components useful in the practice of the subject application include, e.g. cryogenic protectors such as mannitol (to protect from degradation during lyophilization), preservatives, antioxidants, etc. (column 4, lines 45-48). Ron's preferred osteogenic protein is recombinant BMP-2, but any isolated or recombinant osteogenic proteins of the TGF-β family would be similarly useful (column 1, lines 13-34 and 63-65; and column 2, lines 12-14 and 22-24).

Avis teaches that the particular advantages of freeze-drying (lyophilization) are ease of processing a liquid, pharmaceuticals can be stored in a dry state in which there are relatively few stability problems, the products are often more soluble and/or more rapidly soluble, and

Art Unit: 1647

5

10

15

20

dispersions are stabilized throughout their shelf-life (page 1565, paragraph bridging columns 1-2, through column 2, full paragraph 1). Avis teaches that mannitol has been found to be most useful to increase the solids content of the original solution to between approximately 5 and 25% so that the freeze-dried product plug occupies essentially the same volume as that of the original solution (page 1566, column 2, full paragraphs 1-3). A 5 to 25% mannitol solution contains 50 to 250 mg mannitol per ml.

Ron and Avis do not teach lyophilization of MP52 in the presence of mannitol.

However, it would have been obvious to one of ordinary skill in the art at the time of Applicants' invention to make a pharmaceutical composition comprising recombinantly produced MP52, as taught by Neidhardt, and to modify that teaching by making an osteogenic protein in the form of a pharmaceutically acceptable solution (including reconstitution from a lyophilized form), wherein the osteogenic protein is solubilized at a concentration of at least about 2 to 4 mg/ml, said solution further comprising cryogenic protectors such as mannitol, as taught by Ron, with a reasonable expectation of success. One of ordinary skill in the art would be motivated to make this modification because it is optimal to solubilize osteogenic proteins at a concentration of at least about 2 to 4 mg/ml so that a pharmaceutically effective amount of protein can be delivered without undue volumes of carrier being necessary, and because cryogenic protectors, such as mannitol, protect from degradation during lyophilization. It would have been further obvious to one of ordinary skill in the art at the time of Applicants' invention to add mannitol to a purified solution of human MP52 in an amount to give a mixing ratio of 2 to 4 mg/ml MP52 to 50 to 250 mg/ml mannitol and then lyophilize the resultant mixed solution with a reasonable expectation of success. In so doing, one would obtain a solution

Art Unit: 1647

5

10

15

20

comprising MP52 and mannitol, wherein the concentration of mannitol is 0.5-5% (w/v). One of ordinary skill in the art would have been further motivated to make this modification because the particular advantages of freeze-drying (lyophilization) are ease of processing a liquid, pharmaceuticals can be stored in a dry state in which there are relatively few stability problems, the products are often more soluble and/or more rapidly soluble, dispersions are stabilized throughout their shelf-life, and because mannitol has been found to be most useful to increase the solids content of the original solution to between approximately 5 and 25% so that the freeze-dried product plug occupies essentially the same volume as that of the original solution. A solution comprising 2 to 4 mg/ml MP52 and 50 to 250 mg/ml mannitol is a solution comprising MP52 and mannitol in the range of 1 : 5-50 (ratio by weight). A lyophilized form of said solution is a composition comprising MP52 and mannitol in the range of 1 : 5-50 (ratio by weight).

Regarding the limitations "prevention of coloration," "prevention of shrinking," and "prevention of aggregation," *prima facie* obviousness does not require that the prior references suggest combining their disclosure for the same reasons that Applicants combined them. Ron specifically suggest using mannitol as a cryoprotectant to prevent degradation of BMPs during the lyophilization process, which fairly suggest using mannitol during lyophilization of MP52 (a member of the same osteogenic protein family as BMPs). It would have been obvious to one of ordinary skill in the art at the time of Applicants' invention to prevent degradation as much as possible for as long as possible in order to have an as potent as possible for as long as possible MP52 biologic or pharmaceutical, which is motivation to prevent degradation by adding mannitol in accordance with the teachings of Ron. Furthermore, the examiner does not agree

Art Unit: 1647

5

10

15

20

that these results would have been unexpected to one skilled in the art. Discovery of an optimum value of a result effective variable in a known process is ordinarily within the skill of the art. According to Avis, one skilled in the art of lyophilization typically considers "the nature and stability characteristics required during the liquid state, both freshly prepared and when reconstituted before use, [as well as] the characteristics of the dried plug" (Avis, page 1566), when formulating a pharmaceutical or biological product, i.e., "whether the lyophilized substance will be dull or spongy or sparkling and crystalline, firm or friable, expanded or shrunken, etc." (*id.*). It is fair to say that Avis identifies the choice of excipient as a "result effective variable." Identification of mannitol as the ideal lyophilization excipient for MP52 would have been within the ordinary skill of the art.

The invention is prima facie obvious over the prior art.

Claims 7–14 are rejected under 35 U.S.C. 103(a) as being unpatentable over Neidhardt (WO 93/16099) in view of Ron (U. S. Patent No. 5,171,579) and Avis (1990) as applied to claims 7 and 12–14 above and further in view of Chang (J Pharm Sci. 1996 Dec;85(12):1325-30).

Neidhardt in view of Ron and Avis teach a solution comprising MP52 and mannitol and a lyophilized form of said solution, as discussed above. Neidhardt in view of Ron and Avis do not teach a solution comprising MP52 and mannitol and a lyophilized form of said solution, wherein said solution comprises a surfactant.

Chang teaches that the addition of small amounts of surface-active agents protected proteins from both freeze- and surface-induced denaturation (Abstract). Freezing plays a

Art Unit: 1647

5

10

15

20

crucial role in the damage incurred by proteins during freeze-drying (page 1325, left column, full paragraph 1). It has been found with a few proteins that low concentrations of surfactants (i.e., below the critical micellar concentration), which would not be expected to greatly alter the free energy of protein denaturation, provide a high degree of protection during freeze-thawing. Surfactants are known to stabilize proteins against surface-induced denaturation, so these results also suggest that the ice-water interface can contribute to freeze-induced protein denaturation. See page 1325, right column, full paragraph 1. Chang's results indicate that the freezinginduced denaturation is related to the exposure of proteins to an ice-water interface, so it seems rational to use surfactants as cryoprotectants (page 1327, right column, full paragraph 2). The capacity of 0.01% Tween 80 to protect proteins during freeze-thawing appears to be quite general because all of the model proteins were essentially completely protected (paragraph bridging pages 1327-1328). To determine how general this protective effect was, the influence of several surfactants, with different chemical structures, on freeze denaturation of LDH was tested. All the tested surfactants protected LDH from precipitation during a quench-freezing process, even though the control frozen without a surfactant showed a significant increase in turbidity (Table 2). See page 1328, left column, full paragraph 1. The surfactants tested included Tween, Triton, and Brij. See page 1328, Table 2. This general stabilization of proteins during freeze-thawing by relatively low concentrations of surfactants strongly supports the contention that damage to proteins during freezing is due, at least to a large degree, to surface denaturation (page 1328, left column, full paragraph 2). In general, it appears that to obtain native, nonaggregated protein after freeze-drying and rehydration, it is necessary to develop a formulation that protects the protein during both freezing and drying. When 0.1% of

Art Unit: 1647

5

10

15

20

Tween 80 was included in the protein solution, the soluble aggregate content decreased to 3%. Inclusion of 1% sucrose in the formulation, which is known inhibit unfolding of proteins during freeze-drying, led to slightly less protection, as reflected by a 8% aggregate content upon reconstitution. See paragraph bridging pages 1328-1329. To ascertain at what point during the freeze-drying and rehydration process Tween 80 was providing its beneficial effects, Chang conducted the following experiments. First, Chang found that the surfactant can prevent the formation of aggregates during rehydration. When the protein was freeze dried without surfactant but reconstituted with 0.1% Tween 80, the aggregate content was reduced to 23%. However, this protection was not sufficient to explain the reduction of aggregate content to 3% when Tween 80 was included prior to the freeze-drying. See page 1329, left column, full paragraph 1. Thus, for the rational design of stable protein formulations for freezing and freezedrying, it is advantageous to include a surfactant, which apparently inhibits the denaturation of proteins during freezing. A surfactant may only be sufficient to protect proteins during the freezing step. Another stabilizer, which is known to confer protection during drying (e.g., sucrose) will probably be needed to completely inhibit protein unfolding during freeze-drying. See page 1329, right column, last paragraph. Freeze-drying of IL-1ra solutions was performed as follows. An IL-1ra solution containing 2 mM potassium phosphate buffer, 3% mannitol, and 1 or 100 mg/mL of protein was prepared by dialysis. Mannitol was included as a crystalline bulking agent. Addition of 1% (w/v) sucrose or 0.1% (w/v) Tween 80 was made after the dialysis. Each vial was frozen by immersion into liquid nitrogen and loaded onto a shelf that was prechilled to -50 °C. After 1 h at -50 °C, the shelf temperature was increased to -15 °C at a rate of 1 °C/h. After an additional hour of incubation of the vials at this temperature, drying

Art Unit: 1647

5

10

15

20

was initiated by decreasing the chamber pressure to 100 µmHg. Then, the shelf temperature was increased to 25 °C at a rate of 1 °C/min. After completion of a 10-h drying cycle under vacuum the vial head spaces were filled with dry gaseous nitrogen. The vials were capped with stoppers until further analysis. See page 1326, left column, full paragraph 4. It is fair to say that Chang identifies the addition of surfactants and their combination with another stabilizer which is known to confer protection during drying, as a "result effective variable" for inhibiting protein denaturation during freeze-drying. Chang does not teach a lyophilized composition comprising MP52 and mannitol, a process for the preparation of said composition, a solution comprising MP52 and mannitol, or said solution further comprising a detergent/substance.

However, it would have been obvious to one of ordinary skill in the art at the time of Applicants' invention to make teach a solution comprising MP52 and mannitol and a lyophilized form of said solution, as Neidhardt in view of Ron and Avis, and to modify that teaching by adding a surfactant, as taught by Chang, with a reasonable expectation of success. One of ordinary skill in the art would be motivated to make this modification because for the rational design of stable protein formulations for freezing and freeze-drying, it is advantageous to include a surfactant, which apparently inhibits the denaturation of proteins during freezing. The invention is prima facie obvious over the prior art.

Claims 7–10 and 12–14 are rejected under 35 U.S.C. 103(a) as being unpatentable over Neidhardt (WO 93/16099), Ron (U. S. Patent No. 5,171,579), and Avis (1990) as applied to claims 7 and 12–14 above and further in view of Chang (J Pharm Sci. 1996 Dec;85(12):1325-30) and further in view of Hansen (U. S. Patent No. 6,586,574) and in light of the MeSH

Art Unit: 1647

5

10

15

20

definition of "poloxamer."

Neidhardt in view of Ron and Avis and further in view of Chang teach a solution comprising MP52, mannitol, and a surfactant, as discussed above. Neidhardt in view of Ron and Avis and further in view of Chang do not teach a solution comprising MP52, mannitol, and a polyoxyethylene-polyoxypropylene copolymer.

Hansen relates generally to the stabilization of freeze-dried proteins. Hansen discloses that further stabilization of freeze-dried proteins can be obtained by the addition of surfactants, such as Tween and poloxamers (column 6, full paragraph 3). Poloxamers are polyoxyethylene-polyoxypropylene copolymers, as evidenced by the MeSH definition of "poloxamer." The term "surfactants" generally include those agents, which protect the protein from air/solution interface-induced stresses and solution/surface induced-stresses (e.g. resulting in protein aggregation), and may include detergents such as polysorbate, poloxamer or polyethylene glycol, and the like. Optionally, concentrations from about 0.01% to about 1% (w/w) are suitable for maintaining protein stability, however, the levels used in actual practice are customarily limited by clinical practice. See column 5, full paragraph 2. Hansen does not teach a solution comprising MP52, mannitol, and a polyoxyethylene-polyoxypropylene copolymer.

However, it would have been obvious to one of ordinary skill in the art at the time of Applicants' invention to make a solution comprising MP52, mannitol, and a surfactant, as taught by Neidhardt in view of Ron and Avis and further in view of Chang, and to modify that teaching by making a solution comprising MP52, mannitol, and a poloxamer, as taught by Hansen, with a reasonable expectation of success. One of ordinary skill in the art would be motivated to make this modification because further stabilization of freeze-dried proteins can be

Application/Control Number: 10/666,535 Page 11

Art Unit: 1647

obtained by the addition of surfactants, such as poloxamers. The invention is prima facie obvious over the prior art.

Claim Rejections - 35 USC § 112

The following is a quotation of the second paragraph of 35 U.S.C. 112:

The specification shall conclude with one or more claims particularly pointing out and distinctly claiming the subject matter which the applicant regards as his invention.

Claims 8 and 9 are rejected under 35 U.S.C. 112, second paragraph, as being indefinite for failing to particularly point out and distinctly claim the subject matter which applicant regards as the invention.

Claims 8 and 9 are indefinite over the recitation of "detergent/substance" and/or "polyoxyethylenic detergent/substance" because it is unclear if the solution further comprises a detergent and/or a detergent substance or a non-detergent substance, a polyoxyethylenic detergent and/or a polyoxyethylenic detergent substance or a polyoxyethylenic non-detergent substance. The metes and bounds are not clearly set forth.

Conclusion

No claims are allowable.

TO THE GROUP RECEPTIONIST WHOSE TELEPHONE NUMBER IS (703) 308-0196.

ANY INQUIRY CONCERNING THIS COMMUNICATION OR EARLIER COMMUNICATIONS FROM THE EXAMINER SHOULD BE DIRECTED TO DAVID S. ROMEO WHOSE TELEPHONE NUMBER IS (571) 272-0890. THE EXAMINER CAN NORMALLY BE REACHED ON MONDAY THROUGH FRIDAY FROM 7:30 A.M. TO 4:00 P.M. IF ATTEMPTS TO REACH THE EXAMINER BY TELEPHONE ARE UNSUCCESSFUL, THE EXAMINER'S SUPERVISOR, BRENDA BRUMBACK, CAN BE REACHED ON (571) 272-0961.

IF SUBMITTING OFFICIAL CORRESPONDENCE BY FAX, APPLICANTS ARE ENCOURAGED TO SUBMIT OFFICIAL CORRESPONDENCE TO THE CENTRAL FAX NUMBER FOR OFFICIAL CORRESPONDENCE, WHICH IS (571) 273-8300.

CUSTOMERS ARE ALSO ADVISED TO USE CERTIFICATE OF FACSIMILE PROCEDURES WHEN SUBMITTING A REPLY TO A NON-FINAL OR FINAL OFFICE ACTION BY FACSIMILE (SEE 37 CFD 1.6 AND 1.8)

OR FINAL OFFICE ACTION BY FACSIMILE (SEE 37 CFR 1.6 AND 1.8).

ANY INQUIRY OF A GENERAL NATURE OR RELATING TO THE STATUS OF THIS APPLICATION OR PROCEEDING SHOULD BE DIRECTED.

Done Rome

DAVID ROMEO
PRIMARY EXAMINER
ART UNIT 1647

30

25

10

15

35

DSR JUNE 24, 2006